

BIBLIOGRAFÍA

- Abe D, Doi H, Asai T, Kimura M, Wada T, Takahashi Y, et al. Association between COMT Val158Met polymorphism and competition results of competitive swimmers. *J Sports Sci.* 2018;36. doi:10.1080/02640414.2017.1309058
- ACE gene. Available: <https://medlineplus.gov/genetics/gene/ace/>
- Ae CH, Bryan AD, Hagger MS. What keeps a body moving? The brain-derived neurotrophic factor val66met polymorphism and intrinsic motivation to exercise in humans. *J Behav Med.* 2014;37. doi:10.1007/s10865-014-9567-4
- Aleksandra Z, Zbigniew J, Waldemar M, Agata L-D, Mariusz K, Marek S, et al. The Gene M235T Polymorphism and Response of Power-Related Variables to Aerobic Training. *J Sports Sci Med.* 2016;15: 616–624.
- Alves GB, Oliveira EM, Alves CR, Rached HRS, Mota GFA, Pereira AC, et al. Influence of angiotensinogen and angiotensin-converting enzyme polymorphisms on cardiac hypertrophy and improvement on maximal aerobic capacity caused by exercise training. *Eur J Cardiovasc Prev Rehabil.* 2009;16: 487–492.
- Amorim CEG, Acuña-Alonzo V, Salzano FM, Bortolini MC, Hünemeier T. Differing evolutionary histories of the ACTN3*R577X polymorphism among the major human geographic groups. *PLoS One.* 2015;10: e0115449.
- AMPD1 gene. Available: <https://medlineplus.gov/genetics/gene/ampd1/>
- Baumert P, Lake MJ, Stewart CE, Drust B, Erskine RM. Genetic variation and exercise-induced muscle damage: implications for athletic performance, injury and ageing. *Eur J Appl Physiol.* 2016;116: 1595–1625.
- Bell-Pedersen D, Cassone VM, Earnest DJ, Golden SS, Hardin PE, Thomas TL, et al. Circadian rhythms from multiple oscillators: lessons from diverse organisms. *Nat Rev Genet.* 2005;6. doi:10.1038/nrg1633
- Benedetti F, Dallaspezia S, Fulgosi MC, Lorenzi C, Serretti A, Barbini B, et al. Actimetric evidence that CLOCK 3111 T/C SNP influences sleep and activity patterns in patients affected by bipolar depression. *Am J Med Genet B Neuropsychiatr Genet.* 2007;144B. doi:10.1002/ajmg.b.30475
- Benyamin B, Ferreira MAR, Willemsen G, Gordon S, Middelberg RPS, McEvoy BP, et al. Common variants in TMPRSS6 are associated with iron status and erythrocyte volume. *Nat Genet.* 2009;41: 1173.
- Benyamin B, McRae AF, Zhu G, Gordon S, Henders AK, Palotie A, et al. Variants in TF and HFE Explain 40% of Genetic Variation in Serum-Transferrin Levels. *Am J Hum Genet.* 2009;84: 60.
- Bouchard C, Antunes-Correa LM, Ashley EA, Franklin N, Hwang PM, Mattsson CM, et al. Personalized preventive medicine: genetics and the response to regular exercise in preventive interventions. *Prog Cardiovasc Dis.* 2015;57. doi:10.1016/j.pcad.2014.08.005
- Bouillon R, Verstuyf A. Vitamin D, mitochondria, and muscle. *J Clin Endocrinol Metab.* 2013;98. doi:10.1210/jc.2013-1352
- Bryan A, Hutchison KE, Seals DR, Allen DL. A transdisciplinary model integrating genetic, physiological, and psychological correlates of voluntary exercise. *Health Psychol.* 2007;26: 30–39.
- Chard MD, Cawston TE, Riley GP, Gresham GA, Hazleman BL. Rotator cuff degeneration and lateral epicondylitis: a comparative histological study. *Ann Rheum Dis.* 1994;53: 30–34.
- Cieszczyk P, Ostanek M, Leońska-Duniec A, Sawczuk M, Maciejewska A, Eider J, et al. Distribution of the AMPD1 C34T polymorphism in Polish power-oriented athletes. *J Sports Sci.* 2012;30. doi:10.1080/02640414.2011.623710

- COL5A1 gene. Available: <https://medlineplus.gov/genetics/gene/col5a1/>
- Colaboradores de los proyectos Wikimedia. Citoquina. Wikimedia Foundation, Inc.; 6 Jan 2004. Available: <https://es.wikipedia.org/wiki/Citoquina>
- Cole CR, Blackstone EH, Pashkow FJ, Snader CE, Lauer MS. Heart-rate recovery immediately after exercise as a predictor of mortality. *N Engl J Med*. 1999;341. doi:10.1056/NEJM199910283411804
- Cupeiro R, Benito PJ, Maffulli N, Calderón FJ, González-Lamuño D. MCT1 genetic polymorphism influence in high intensity circuit training: a pilot study. *J Sci Med Sport*. 2010;13: 526–530.
- Cupeiro R, González-Lamuño D, Amigo T, Peinado AB, Ruiz JR, Ortega FB, et al. Influence of the MCT1-T1470A polymorphism (rs1049434) on blood lactate accumulation during different circuit weight trainings in men and women. *J Sci Med Sport*. 2012;15: 541–547.
- Del Coso J, Hiam D, Houweling P, Pérez LM, Eynon N, Lucía A. More than a “speed gene”: ACTN3 R577X genotype, trainability, muscle damage, and the risk for injuries. *Eur J Appl Physiol*. 2018;119: 49–60.
- El Khoury L, Ribbans WJ, Raleigh SM. MMP3 and TIMP2 gene variants as predisposing factors for Achilles tendon pathologies: Attempted replication study in a British case-control cohort. *Meta Gene*. 2016;9: 52–55.
- Evangelou E, Chapman K, Meulenbelt I, Karassa FB, Loughlin J, Carr A, et al. Large-scale analysis of association between GDF5 and FRZB variants and osteoarthritis of the hip, knee, and hand. *Arthritis Rheum*. 2009;60. doi:10.1002/art.24524
- Evans DM, Zhu G, Dy V, Heath AC, Madden PAF, Kemp JP, et al. Genome-wide association study identifies loci affecting blood copper, selenium and zinc. *Hum Mol Genet*. 2013;22: 3998.
- Ferguson BS, Rogatzki MJ, Goodwin ML, Kane DA, Rightmire Z, Gladden LB. Lactate metabolism: historical context, prior misinterpretations, and current understanding. *Eur J Appl Physiol*. 2018;118. doi:10.1007/s00421-017-3795-6
- Galasso O, Iaccino E, Gallelli L, Perrotta I, Conforti F, Donato G, et al. Collagen type V polymorphism in spontaneous quadriceps tendon ruptures. *Orthopedics*. 2012;35. doi:10.3928/01477447-20120327-31
- Garton FC, Houweling PJ, Vukcevic D, Meehan LR, Lee FXZ, Lek M, et al. The Effect of ACTN3 Gene Doping on Skeletal Muscle Performance. *Am J Hum Genet*. 2018;102: 845.
- Genetic profile of elite endurance athletes. *Sports, Exercise, and Nutritional Genomics*. Academic Press; 2019. pp. 73–104.
- Genetics and Sprint, Strength, and Power Performance: Candidate Gene Versus Genome-Wide Association Study Approaches. *Nutrition and Enhanced Sports Performance*. Academic Press; 2019. pp. 371–383.
- Girgis CM, Clifton-Bligh RJ, Hamrick MW, Holick MF, Gunton JE. The roles of vitamin D in skeletal muscle: form, function, and metabolism. *Endocr Rev*. 2013;34. doi:10.1210/er.2012-1012
- Girgis CM, Mokbel N, Cha KM, Houweling PJ, Abboud M, Fraser DR, et al. The Vitamin D Receptor (VDR) Is Expressed in Skeletal Muscle of Male Mice and Modulates 25-Hydroxyvitamin D (25OHD) Uptake in Myofibers. *Endocrinology*. 2014;155: 3227.
- Gomez-Gallego F, Santiago C, González-Freire M, Yvert T, Muniesa CA, Serratos L, et al. The C allele of the AGT Met235Thr polymorphism is associated with power sports performance. *Appl Physiol Nutr Metab*. 2009;34. doi:10.1139/H09-108
- Graham TE. Caffeine and exercise: metabolism, endurance and performance. *Sports Med*. 2001;31. doi:10.2165/00007256-200131110-00002
- Guest N, Corey P, Vescovi J, El-Sohemy A. Caffeine, CYP1A2 Genotype, and Endurance Performance in Athletes. *Med Sci Sports Exerc*. 2018;50: 1570–1578.

- Guest NS, VanDusseldorp TA, Nelson MT, Grgic J, Schoenfeld BJ, Jenkins NDM, et al. International society of sports nutrition position stand: caffeine and exercise performance. *J Int Soc Sports Nutr.* 2021;18: 1–37.
- Gureev AP, Shaforostova EA, Popov VN. Regulation of Mitochondrial Biogenesis as a Way for Active Longevity: Interaction Between the Nrf2 and PGC-1 α Signaling Pathways. *Front Genet.* 2019;10. doi:10.3389/fgene.2019.00435
- Hautala AJ, Rankinen T, Kiviniemi AM, Mäkikallio TH, Huikuri HV, Bouchard C, et al. Heart rate recovery after maximal exercise is associated with acetylcholine receptor M2 (CHRM2) gene polymorphism. *Am J Physiol Heart Circ Physiol.* 2006;291: H459–66.
- Hautala AJ, Tulppo MP, Kiviniemi AM, Rankinen T, Bouchard C, Mäkikallio TH, et al. Acetylcholine receptor M2 gene variants, heart rate recovery, and risk of cardiac death after an acute myocardial infarction. *Ann Med.* 2009;41. doi:10.1080/07853890802477866
- Jones A, Montgomery HE, Woods DR. Human performance: a role for the ACE genotype? *Exerc Sport Sci Rev.* 2002;30. doi:10.1097/00003677-200210000-00008
- Khoschnau S, Melhus H, Jacobson A, Rahme H, Bengtsson H, Ribom E, et al. Type I collagen alpha1 Sp1 polymorphism and the risk of cruciate ligament ruptures or shoulder dislocations. *Am J Sports Med.* 2008;36. doi:10.1177/0363546508320805
- Kiah McCabe CC. Can Genetics Predict Sports Injury? The Association of the Genes GDF5, AMPD1, COL5A1 and IGF2 on Soccer Player Injury Occurrence. *Sportscience.* 2018;6. doi:10.3390/sports6010021
- Kim S, Kleimyer JP, Ahmed MA, Avins AL, Fredericson M, Dragoo JL, et al. A Genetic Marker Associated with Shoulder Dislocation. *Int J Sports Med.* 2017;38. doi:10.1055/s-0043-106190
- Kozlovskaja M, Vlahovich N, Ashton KJ, Hughes DC. Biomedical Risk Factors of Achilles Tendinopathy in Physically Active People: a Systematic Review. *Sports Med Open.* 2017;3: 20.
- Lemme NJ, Li NY, DeFroda SF, Kleiner J, Owens BD. Epidemiology of Achilles Tendon Ruptures in the United States: Athletic and Nonathletic Injuries From 2012 to 2016. *Orthopaedic Journal of Sports Medicine.* 2018;6. doi:10.1177/2325967118808238
- Lopez-Leon S, Tuvblad C, Forero DA. Sports genetics: the PPAR α gene and athletes' high ability in endurance sports. A systematic review and meta-analysis. *Biol Sport.* 2016;33: 3.
- Lortie G, Simoneau JA, Hamel P, Boulay MR, Landry F, Bouchard C. Responses of maximal aerobic power and capacity to aerobic training. *Int J Sports Med.* 1984;5. doi:10.1055/s-2008-1025911
- Mann TN, Lamberts RP, Lambert MI. High responders and low responders: factors associated with individual variation in response to standardized training. *Sports Med.* 2014;44. doi:10.1007/s40279-014-0197-3
- Massidda M, Eynon N, Bachis V, Corrias L, Culigioni C, Piras F, et al. Influence of the MCT1 rs1049434 on Indirect Muscle Disorders/Injuries in Elite Football Players. *Sports Med Open.* 2015;1: 33.
- Massidda M, Mendez-Villanueva A, Ginevičienė V, Proia P, Drozdovska SB, Dosenko V, et al. Association of Monocarboxylate Transporter-1 (MCT1) A1470T Polymorphism (rs1049434) with Forward Football Player Status. *Int J Sports Med.* 2018;39. doi:10.1055/a-0634-6387
- Miyamoto Y, Mabuchi A, Shi D, Kubo T, Takatori Y, Saito S, et al. A functional polymorphism in the 5' UTR of GDF5 is associated with susceptibility to osteoarthritis. *Nat Genet.* 2007;39. doi:10.1038/2005
- Molecular mechanism of PPAR α action and its impact on lipid metabolism, inflammation and fibrosis in non-alcoholic fatty liver disease. *J Hepatol.* 2015;62: 720–733.
- Moore GE, Shuldiner AR, Zmuda JM, Ferrell RE, McCole SD, Hagberg JM. Obesity gene variant and elite endurance performance. *Metabolism.* 2001;50. doi:10.1053/meta.2001.28140
- Moreau ME, Garbacki N, Molinaro G, Brown NJ, Marceau F, Adam A. The kallikrein-kinin system:

current and future pharmacological targets. *J Pharmacol Sci.* 2005;99. doi:10.1254/jphs.srj05001x

• Mori M, Higuchi K, Sakurai A, Tabara Y, Miki T, Nose H. Genetic basis of inter-individual variability in the effects of exercise on the alleviation of lifestyle-related diseases. *J Physiol.* 2009;587. doi:10.1113/jphysiol.2009.179283

• Morici G, Gruttad'Auria CI, Baiamonte P, Mazzuca E, Castrogiovanni A, Bonsignore MR. Endurance training: is it bad for you? *Breathe.* 2016;12: 140.

• Nishime EO, Cole CR, Blackstone EH, Pashkow FJ, Lauer MS. Heart Rate Recovery and Treadmill Exercise Score as Predictors of Mortality in Patients Referred for Exercise ECG. *JAMA.* 2000;284: 1392–1398.

• Norman B, Esbjörnsson M, Rundqvist H, Osterlund T, von Walden F, Tesch PA. Strength, power, fiber types, and mRNA expression in trained men and women with different ACTN3 R577X genotypes. *J Appl Physiol.* 2009;106: 959–965.

• North KN, Yang N, Wattanasirichaigoon D, Mills M, Eastel S, Beggs AH. A common nonsense mutation results in alpha-actinin-3 deficiency in the general population. *Nat Genet.* 1999;21: 353–354.

• O'Connell K, Posthumus M, Schwellnus MP, Collins M. Collagen genes and exercise-associated muscle cramping. *Clin J Sport Med.* 2013;23. doi:10.1097/JSM.0b013e3182686aa7

• Pickering C, Kiely J, Suraci B, Collins D. The magnitude of Yo-Yo test improvements following an aerobic training intervention are associated with total genotype score. *PLoS One.* 2018;13. doi:10.1371/journal.pone.0207597

• Pickering C, Kiely J. Are the Current Guidelines on Caffeine Use in Sport Optimal for Everyone? Inter-individual Variation in Caffeine Ergogenicity, and a Move Towards Personalised Sports Nutrition. *Sports Med.* 2018;48: 7–16.

• Posthumus M, Collins M, Cook J, Handley CJ, Ribbans WJ, Smith RK, et al. Components of the transforming growth factor-beta family and the pathogenesis of human Achilles tendon pathology--a genetic association study. *Rheumatology.* 2010;49. doi:10.1093/rheumatology/keq072

• Prior SJ, Hagberg JM, Phares DA, Brown, Fairfull L, Ferrell RE, et al. Sequence variation in hypoxia-inducible factor 1alpha (HIF1A): association with maximal oxygen consumption. *Physiol Genomics.* 2003;15. doi:10.1152/physiolgenomics.00061.2003

• Raleigh SM, van der Merwe L, Ribbans WJ, Smith RK, Schwellnus MP, Collins M. Variants within the MMP3 gene are associated with Achilles tendinopathy: possible interaction with the COL5A1 gene. *Br J Sports Med.* 2009;43. doi:10.1136/bjism.2008.053892

• Roth SM. Genetic aspects of skeletal muscle strength and mass with relevance to sarcopenia. *BoneKey Reports.* 2012;1. doi:10.1038/bonekey.2012.58

• Ruiz JR, Buxens A, Artieda M, Arteta D, Santiago C, Rodríguez-Romo G, et al. The -174 G/C polymorphism of the IL6 gene is associated with elite power performance. *J Sci Med Sport.* 2010;13. doi:10.1016/j.jsams.2009.09.005

• Ryan RM, Deci EL. Self-determination theory and the facilitation of intrinsic motivation, social development, and well-being. *Am Psychol.* 2000;55. doi:10.1037//0003-066x.55.1.68

• Sarpeshkar V, Bentley DJ. Adrenergic-β 2 receptor polymorphism and athletic performance. *J Hum Genet.* 2010;55: 479–485.

• Saunders CJ, van der Merwe L, Posthumus M, Cook J, Handley CJ, Collins M, et al. Investigation of variants within the COL27A1 and TNC genes and Achilles tendinopathy in two populations. *J Orthop Res.* 2013;31: 632–637.

• SelfDecode Available: <https://selfdecode.com/blog/article/il6-athletic-performance-68/>

• Serrano JM, Banks JB, Fagan TJ, Tartar JL. The influence of Val158Met COMT on physiological stress responsivity. *Stress.* 2019;22. doi:10.1080/10253890.2018.1553949

- Snyder EM, Johnson BD, Joyner MJ. Genetics of β 2-Adrenergic Receptors and the Cardiopulmonary Response to Exercise. *Exerc Sport Sci Rev.* 2008;36: 98.
- Southam L, Rodriguez-Lopez J, Wilkins JM, Pombo-Suarez M, Snelling S, Gomez-Reino JJ, et al. An SNP in the 5'-UTR of GDF5 is associated with osteoarthritis susceptibility in Europeans and with in vivo differences in allelic expression in articular cartilage. *Hum Mol Genet.* 2007;16. doi:10.1093/hmg/ddm174
- Stefan N, Thamer C, Staiger H, Machicao F, Machann J, Schick F, et al. Genetic variations in PPARG and PPARGC1A determine mitochondrial function and change in aerobic physical fitness and insulin sensitivity during lifestyle intervention. *J Clin Endocrinol Metab.* 2007;92. doi:10.1210/jc.2006-1785
- Steinbacher P, Feichtinger RG, Kedenko L, Kedenko I, Reinhardt S, Schönauer A-L, et al. The Single Nucleotide Polymorphism Gly482Ser in the PGC-1 α Gene Impairs Exercise-Induced Slow-Twitch Muscle Fibre Transformation in Humans. *PLoS One.* 2015;10: e0123881.
- Tharabenjasin P, Pabalan N, Jarjanazi H. Association of the ACTN3 R577X (rs1815739) polymorphism with elite power sports: A meta-analysis. *PLoS One.* 2019;14. doi:10.1371/journal.pone.0217390
- Valdivieso P, Vaughan D, Laczko E, Brogioli M, Waldron S, Rittweger J, et al. The Metabolic Response of Skeletal Muscle to Endurance Exercise Is Modified by the ACE-I/D Gene Polymorphism and Training State. *Front Physiol.* 2017;8. doi:10.3389/fphys.2017.00993
- Valeeva EV, Kashevarov GS, Kasimova RR, Ahmetov II, Kravtsova OA. Association of the Val158Met Polymorphism of the COMT Gene with Measures of Psychophysiological Status in Athletes. *Neurosci Behav Physiol.* 2020;50: 485–492.
- Varley I, Greeves JP, Sale C, Friedman E, Moran DS, Yanovich R, et al. Functional polymorphisms in the P2X7 receptor gene are associated with stress fracture injury. *Purinergic Signal.* 2016;12. doi:10.1007/s11302-016-9495-6
- Wagner H, Thaller S, Dahse R, Sust M. Biomechanical muscle properties and angiotensin-converting enzyme gene polymorphism: a model-based study. *Eur J Appl Physiol.* 2006;98. doi:10.1007/s00421-006-0293-7
- Wolfarth B, Rankinen T, Mühlbauer S, Scherr J, Boulay MR, Pérusse L, et al. Association between a beta2-adrenergic receptor polymorphism and elite endurance performance. *Metabolism.* 2007;56. doi:10.1016/j.metabol.2007.07.006
- Yamin C, Duarte JAR, Oliveira JMF, Amir O, Sagiv M, Eynon N, et al. IL6 (-174) and TNFA (-308) promoter polymorphisms are associated with systemic creatine kinase response to eccentric exercise. *Eur J Appl Physiol.* 2008;104: 579–586.
- Zhang B, Tanaka H, Shono N, Miura S, Kiyonaga A, Shindo M, et al. The I allele of the angiotensin-converting enzyme gene is associated with an increased percentage of slow-twitch type I fibers in human skeletal muscle. *Clin Genet.* 2003;63: 139–144.
- Zhang L, Ptáček LJ, Fu Y-H. Diversity of Human Clock Genotypes and Consequences. *Prog Mol Biol Transl Sci.* 2013;119: 51